



Keeping
Your Liver
Healthy



Hepatitis Basics

A Basic Guide to Hepatitis B

Alan Franciscus

Executive Director
Editor-in-Chief, HCSP Publications

The information in this guide is designed to help you understand and manage HBV and is not intended as medical advice. All persons with HBV should consult a medical practitioner for diagnosis and treatment of HBV.



Table of Contents

1. Introduction2
2. Hepatitis B Facts.2
3. Your Liver and Hepatitis2
4. HBV Transmission and Prevention3
5. HBV Prevention Guidelines4
6. The HBV Vaccine and Post-Exposure Prophylaxis4
7. HBV Disease Progression Acute and Chronic HBV4
8. HBV Diagnosis and Monitoring6
9. HBV Treatment.7
10. Managing Drug Side Effects.8
11. HBV Management9
12. Healthy Liver Tips.9
13. Conclusion	11
14. Hepatitis Resources.	11

www.hbvadvocate.org



1. INTRODUCTION

HBV is a blood-borne virus. It enters the body through direct blood exposure and through sexual contact. Fewer than 5% of adults infected with HBV become carriers and remain infected with HBV for longer than six months. For people who are infected with HBV at birth or as young children, the risk of becoming a carrier may be as high as 90%.

An estimated 240 million people worldwide are chronically infected with HBV and there are 786,000 annual deaths. HBV is responsible for up to 80% of all cases of liver cancer worldwide. In 2015, the U.S. Centers for Disease Control and Prevention (CDC) estimated that 19,200 Americans were newly infected with HBV and up to 2.2 million persons have chronic HBV infection. Asians and immigrants from other areas of the world where hepatitis B is endemic have higher rates of infection; most are infected at birth or as young children.

Most people infected with HBV do not have symptoms and lead normal lives. However, in about 20-30% of cases HBV can cause serious liver damage – including fibrosis and cirrhosis – usually over several years or decades. In severe cases, hepatitis B can lead to liver failure and death. An effective vaccine is available to prevent HBV. There are also various treatments that can help slow or stop disease progression.

An estimated 240 million people worldwide are chronically infected with HBV and there are 786,000 annual deaths.

2. HEPATITIS B FACTS

- The CDC estimates that up to 2.2 million Americans are chronically infected with HBV.
- An estimated 1,800 Americans die each year due to complications related to hepatitis B.
- Immigrants from Asia, the Middle East, and parts of Africa—and their children—are at higher risk for HBV infection.

Universal hepatitis B vaccination is recommended by the World Health Organization; today, teenagers are routinely vaccinated against HBV, but infant vaccination is still optional in several states. The American Academy of Pediatrics recommends HBV vaccination for all infants.

3. YOUR LIVER AND HEPATITIS

The liver is the largest internal organ. It is located behind the ribcage on the right side of the abdomen. The liver is responsible for some 500 bodily functions. It processes almost everything you eat, breathe, or absorb through the skin. Liver cells produce bile, which enables the body to digest food and absorb nutrients. The liver plays an important role in metabolism, including the storage of vitamins, minerals, and sugars. It synthesizes several important proteins, including hormones, blood proteins, and clotting factors. The liver also plays a role in detoxification, filtering drugs and toxins (poisons) and eliminating toxic byproducts of normal metabolism.

Hepatitis is a general name for inflammation of the liver. It may be caused by viruses, toxic chemicals, alcohol, drugs, or other factors. HBV attacks and replicates in liver cells. As liver damage progresses, the organ may no longer be able to carry out its normal activities. Because the liver is responsible for so many important functions, liver disease can lead to a variety of different symptoms and conditions.

Hepatitis B is one of the three most common types of viral hepatitis; the others are hepatitis A and hepatitis C. Hepatitis A is most common. The hepatitis A virus (HAV)



is usually transmitted through food or water contaminated by HAV-infected feces, but may also be contracted through household and sexual (anal/oral) contact. There is no treatment for hepatitis A, but the body usually clears the virus and it does not cause chronic infection. Hepatitis C (formerly known as non-A/non-B hepatitis) is a blood-borne virus like hepatitis B. HCV is much more likely than HBV to become chronic, and also is more likely to lead to liver damage. Although the long-term effects of hepatitis B and hepatitis C are similar in many respects, the diseases are caused by two distinct viruses that are not closely related. Both hepatitis A and hepatitis B can be prevented by vaccines; there is currently no vaccine for hepatitis C.

4. HBV TRANSMISSION & PREVENTION


HBV is transmitted by direct blood-to-blood contact. The hepatitis B virus can live up to 7 days outside of the body. A common transmission route is sharing drug equipment for both injection and non-injection drugs (including needles, cookers, tourniquets, cocaine straws, and crack pipes). Needles used for tattooing and body piercing may also spread the virus. Sharing personal items such as razors, toothbrushes, and nail files is a less likely but still possible transmission route. This can happen when a small amount of HBV-infected blood remains on an item after use and is transferred to another person who uses the same item.

In the past, many people contracted HBV through blood transfusions; however, a test for HBV in donated blood has been in use since 1972 and a test for HCV became available in 1992. Today, blood transfusions are considered safe. Health-care workers may be exposed to HBV through needle-sticks and other accidental exposures on the job.

HBV is present in semen and vaginal fluids, and hepatitis B may be transmitted through sexual activity. HBV is much more likely than HCV to be sexually transmitted.

The CDC estimates that the majority of new HBV infections in the United States are sexually transmitted. Transmission may be more likely during a woman's menstrual period. HBV transmission rates are particularly high among men who have sex with men. Perinatal transmission from HBV-infected mothers to their infants before or during birth account for the majority of infections in areas where HBV is endemic. Transmission is more likely if the mother has a high level of HBV in her blood; mothers coinfecting with HCV or HIV in addition to HBV also appear to be more likely to transmit hepatitis B to their babies. Although hepatitis B surface antigen, a particle of the virus, is present in breast milk, there is no evidence that hepatitis B is transmitted through breast-feeding if the infant is vaccinated.

Studies indicate that HBV transmission is common between young children in areas where the virus is endemic, probably due to scratching and biting. Although HBV is detectable in saliva, hepatitis B is not known to be transmitted by sneezing, coughing, or sharing eating



"In the past, many people contracted HBV through blood transfusions; however, a test for HBV in donated blood has been in use since 1972 and a test for HCV became available in 1992."



utensils or drinking glasses; household transmission of HBV rarely occurs. There are no documented cases of HBV transmission through urine, feces, sweat, tears, or vomit. In one-third or more cases, people have no identifiable risk factors and the route of hepatitis B transmission is unknown.

5. HBV PREVENTION GUIDELINES

- Get vaccinated against hepatitis B.
- Infants born to HBV-infected mothers should receive the hepatitis B vaccine and HBV immune globulin (HBIG) on the day of birth.
- Practice safer sex, including the use of condoms and barriers.
- Do not share needles to inject drugs; obtain new needles from a needle exchange or – if you must share – clean needles thoroughly with bleach.
- Do not share drug equipment such as cocaine straws or crack pipes.
- Tattooists, piercers, and acupuncturists should use new needles for each client.
- Manicurists and barbers should disinfect tools between customers or use disposable, single-use items.
- Do not share personal items such as razors, toothbrushes, nail files, or pierced earrings.
- Exercise universal precautions in health-care settings, including the use of safety gloves.
- Properly dispose of used needles, bandages, and menstrual supplies; clean and disinfect spilled body fluids.
- Cover all cuts, sores, and rashes.

6. THE HBV VACCINE AND POST-EXPOSURE PROPHYLAXIS

Hepatitis B can be prevented with a vaccine. The vaccine is administered as a series of three injections given over six months (the second injection one month after the first,

and the third injection five months later). A two-dose vaccine has been approved for adolescents aged 11-15.

A combination HAV/HBV vaccine (Twinrix) is also available. The Food and Drug Administration has approved accelerated dosing of Twinrix (3 shots within 30 days followed by a booster in one year).

The HBV vaccine is also recommended for health care workers, sexually active adults, injection drug users, household contacts of HBV-infected individuals, people infected with HCV, and adolescents who were not vaccinated as children.

There are no documented cases of HBV transmission through urine, feces, sweat, tears, or vomit.

If a person has recently been exposed to HBV, post exposure prophylaxis using the HBV vaccine plus injected antibodies (HBV immune globulin, or HBIG) can help prevent the development of hepatitis B or reduce the length and severity of illness. This procedure should be done within 72 hours after exposure. HBIG plus the HBV vaccine can also prevent hepatitis B in infants born to HBV-positive mothers.

7. HBV DISEASE PROGRESSION ACUTE AND CHRONIC HBV

After exposure to HBV, the incubation period usually lasts 30-90 days. The initial phase of hepatitis B is called acute infection. Clearance of HBV following an acute infection usually takes 2-12 months, during which time a person may experience fatigue and abdominal tenderness.


In a majority of people infected with HBV, the immune system can clear the virus. But some HBV-infected adults



– estimated at less than 6% – will become chronically infected, meaning the virus remains in the body after six months. Among people infected with HBV as infants or children, this number is much higher – up to 90% for infants and 30% for children ages 1-5. HBV genetic material (DNA) remains in the nuclei of the cells of everyone infected with the virus, even when HBV infection cannot be detected in the blood. As a result, HBV may reactivate if people’s immune systems are damaged or if they receive immunosuppressive drugs such as steroids or chemotherapy. A majority (75%) of people with chronic hepatitis B have no symptoms of liver disease, but this situation can change at any time during the life of a carrier.

Hepatitis B Symptoms

Most people with HBV experience few or no symptoms; in fact, many are unaware that they carry the virus. Many people with acute hepatitis B have no symptoms, and most people with chronic HBV also have few or no symptoms. If they do occur, symptoms of acute hepatitis B may include fatigue (unusual, prolonged tiredness), fever, malaise (a flulike feeling), nausea, vomiting, loss of appetite (anorexia), abdominal pain or bloating, indigestion, headaches, itching (pruritus), and muscle or joint aches. Rarely, HBV may be associated with rheumatological problems such as polyarteritis nodosa.



“Cirrhosis – a process in which liver cells are destroyed and replaced with scar tissue. Extensive scar formation can impair the flow of blood through the liver.”

Some people with either acute or chronic hepatitis B may develop jaundice (yellowing of the skin and whites of the eyes), dark urine, and pale-colored stools, caused by a high level of bilirubin (a pigment) in the body. Some people also develop high levels of certain liver enzymes, especially ALT.

Late-Stage Disease

In a minority of people with hepatitis B, the disease progresses over years or decades, with increasing liver damage. An estimated 20-30% of people with chronic HBV will develop cirrhosis. In severe cases, a person may experience liver failure and require a liver transplant.

Liver damage may include:

Inflammation – an immune response to infection or injury characterized by infiltration of white blood cells, swelling, and functional impairment of liver cells. People with liver inflammation may – but do not always – have elevated liver enzyme levels.

Necrosis – the death of liver cells (hepatocytes).

Fibrosis – the development of scar tissue within the liver which, if extensive, may begin to interfere with the smooth flow of blood through the liver.

Cirrhosis – a process in which liver cells are destroyed and replaced with scar tissue. Extensive scar formation can impair the flow of blood through the liver. Compensated cirrhosis is when the liver is scarred but can still work relatively normally; people with compensated cirrhosis usually exhibit few symptoms. Decompensated cirrhosis is when the liver is so damaged that it cannot function properly. People with decompensated cirrhosis may develop complications such as bleeding varices (ruptured blood vessels in the esophagus, stomach, and the gastrointestinal system), abdominal fluid accumulation (ascites), easy bleeding or bruising, mental impairment (hepatic encephalopathy), and coma.



Hepatocellular carcinoma – a type of liver cancer that may occur in people with chronic hepatitis. Liver cancer typically occurs in people with cirrhosis, but some individuals with hepatitis B who develop liver cancer do not have cirrhosis.

8. HBV DIAGNOSIS AND MONITORING

Various tests are used to diagnose hepatitis B and to assess the stage of disease and the extent of liver damage.

Antibody Tests

Hepatitis B is diagnosed and staged by looking at a complex combination of HBV antigens and antibodies. Tests are available to measure three HBV-associated proteins or antigens: HBsAg (surface), HBcAg (core), and HBeAg. The immune system produces three corresponding antibodies: called anti-HBs, anti-HBc, and anti-HBe. The presence of HBsAg and/or HBV DNA in the blood indicates that a person currently has hepatitis B. The presence of anti-HBs antibodies when HBsAg is absent shows that a person does not have active disease. People who have been exposed to HBV and have successfully cleared the infection test positive for both anti-HBs and anti-HBc. People who have received the HBV vaccine have anti-HBs but not anti-HBc.

The presence of HBeAg often indicates that the virus is actively replicating and that a person is highly infectious and at greater risk for liver damage. Traditionally, the loss of HBeAg has been used as an indication that treatment is effective. However, when people have had hepatitis B for many years, they may test negative for HBeAg, but still have an active infection and high viral load. This is called HBeAg-negative hepatitis B. These cases result when HBV mutate and are able to replicate without HBeAg.

Hepatitis B Serology*

Hepatitis B surface antigen (HBsAg): A protein on the surface of HBV; it can be detected in serum (blood) during

acute or chronic HBV infection. The presence of HBsAg indicates a person is infectious. The body normally produces antibodies to HBsAg (anti-HBs) as part of its immune response to fight the infection. HBsAg is the antigen used to make hepatitis B vaccine.

Hepatitis B surface antibody (anti-HBs): The presence of anti-HBs are generally interpreted as indicating recovery and immunity from HBV infection. Anti-HBs also develop in people who have been successfully vaccinated against hepatitis B.

Total hepatitis B core antibody (anti-HBc): Positivity indicates a recent or current infection with HBV (≤ 6 months).

Hepatitis B “e” antigen (HBeAg): This antigen, which can make up part of the virus’s core, is found in serum during acute and chronic hepatitis B. Its presence can indicate that the virus is replicating rapidly and that the person has high levels of HBV. However, it is possible to be infected and have a high viral load if this antigen is absent. This can happen in adults who have had hepatitis B infection for many years. This is called HBeAg-negative hepatitis B.

Hepatitis B “e” antibody (HBeAb or anti-HBe): This antibody is produced by the immune system during acute HBV infection or during active viral replication. Spontaneous conversion from HBeAg to anti-HBe (known as seroconversion) can indicate lower viral load in patients undergoing antiviral or interferon treatment.

**adapted from <https://www.cdc.gov/hepatitis/hbv/hbvfaq.htm#general>*

Viral Load Tests

Viral load tests measure the amount of HBV DNA (genetic material) circulating in the blood. A detectable viral load indicates that HBV is actively replicating. In people with abnormal liver enzyme levels, higher HBV DNA viral load appears to be associated with more severe liver disease. Viral load tests are also useful as an indication of how well antiviral treatment is working.



Biochemical Liver Tests

Biochemical liver tests are a rough indication of the degree of liver inflammation. A hepatic panel includes measurements of various substances in the blood. Many – but not all – people with acute or chronic hepatitis B develop elevated levels of two liver enzymes called alanine aminotransferase (ALT, formerly known as SPGT) and aspartate aminotransferase (AST, formerly known as SGOT). ALT and AST are released into the blood when the liver is damaged. Elevated liver enzyme levels are often the first sign of liver problems, and a decrease in ALT often indicates that treatment is working. However, many people with hepatitis B have persistently normal liver enzyme levels. In addition, some people have normal ALT levels even though they have background cirrhosis.

Another common measure is bilirubin level. Bilirubin is a pigment that is continuously produced as a byproduct of the natural breakdown of red blood cells. A high bilirubin level causes jaundice. Bilirubin level indicates the degree of liver function, as do serum albumin level and measures of blood clotting. Because test results can vary from lab to lab, it is recommended that the same laboratory be used each time so that results can be compared. Keep copies of your lab results for future reference.

HBV Genotypes

There have been 10 HBV genotypes (lettered A through J) or strains identified globally. Furthermore, some HBV genotypes have sub-genotypes. Testing for HBV genotype is important. Different genotypes and sub-genotypes are known to affect the rate of chronicity, disease progression (liver cirrhosis and liver cancer development), treatment choices and treatment outcome. Additionally, certain genotypes and sub-genotypes are more or less prevalent in different regions of the world.

Liver Biopsy

Biopsies are done to assess the extent of inflammation and the amount of scarring of the liver. Biopsy is

the most reliable indicator of liver damage and is used to help make decisions about treatment. Many doctors do not recommend biopsies for HBV carriers with normal biochemical liver tests. In the most common biopsy procedure, the abdominal skin and muscle are numbed and a long, thin needle is quickly inserted into the liver to draw out a small sample of tissue, which is examined under a microscope. Complications from liver biopsies are rare. If you are anxious about the procedure, ask your doctor for a mild tranquilizer prior to the biopsy and for pain medication afterwards.

9. HBV TREATMENT

There are currently two types of drug treatments for chronic hepatitis B: pegylated interferon and antivirals.

Hepatitis B treatment is more likely to be beneficial if a person has elevated liver enzymes (ALT), and low viral load. Treatment for people with very low HBV DNA levels and normal ALT levels is not usually recommended because this shows there is no liver damage occurring. A majority of people chronically infected with HBV will never require treatment, but should have their ALT levels and overall health monitored regularly at least every six months.

There is no cure for HBV. The goal of HBV treatment is to suppress HBV viral load, and HBV disease progression (cirrhosis, liver and death).



Complications from liver biopsies are rare. If you are anxious about the procedure, ask your doctor for a mild tranquilizer prior to the biopsy and for pain medication afterwards.



There are six HBV therapies approved by the Food and Drug Administration (FDA) for the treatment of adults with chronic HBV and five for children with chronic HBV. The American Association for the Study of Liver Disease (AASLD) provides guidelines for the treatment of adults and children with HBV. The links are listed at the end of this Guide.

Alternative Therapies

In addition to pharmaceutical drugs, some people use alternative and complementary therapies for hepatitis B. Herbal remedies should be treated like drugs, since they may have side effects and can interact with other herbs and conventional medications. Many herbs can be toxic to the liver, including chaparral, germander, kava kava, and plants that contain pyrrolizidine alkaloids. Herbs are discussed in our Herbal Glossary at <http://hcvadvocate.org/resources/glossaries/herbal-glossary-2/>. Make sure to contact a reputable herbalist or nutritionist and inform all your health-care providers about any herbs, supplements, or other alternative therapies you are using.

Clinical Trials

There are many medications that are being tested to treat and possibly cure chronic HBV. The process of testing a new drug involves establishing its safety and toxicity in humans (Phase I trials), measuring its safety and tolerability (Phase II trials), and comparing the new drug to current standard treatments or placebo (Phase III trials). After the FDA has granted approval and the new drug is marketed, ongoing studies are done to refine the treatment for maximum safety and effectiveness (Phase IV, or post-marketing trials).

Clinical trials can be a good way to obtain free medication; some trials also may cover the costs of laboratory tests (although they may not provide the results to you or your doctor). Be aware that clinical trial participants typically are randomized to receive either the experimental drug or a standard treatment or placebo (an inactive substance used as a control); you may not be chosen to receive the new drug or the most effective dosage. Read

all the clinical trial information and make sure that you fully understand the terms and conditions before you give your informed consent to participate.

10. MANAGING DRUG SIDE EFFECTS

Drugs used to treat HBV are associated with side effects. Common side effects of pegylated interferon include flulike symptoms, fatigue, nausea, diarrhea, headache, muscle and joint aches, depression, and anxiety; less common side effects include hair loss and thyroid dysfunction. Side effects of antivirals are uncommon, but may include nausea, diarrhea, and headache; rarer side effects include low blood cell counts, peripheral neuropathy (nerve damage in the hands and feet), and pancreatitis (inflammation of the pancreas).



“Side effects of antivirals are uncommon, but may include nausea, diarrhea, and headache.”

Some physical symptoms may be reduced with low dose ibuprofen (do not take if you have cirrhosis) or acetaminophen (high doses can be toxic to the liver). People experiencing psychological symptoms such as depression, anxiety, or irritability may find antidepressants helpful. Some people find that injecting interferon just before bedtime lets them sleep through the worst side effects. Vary where you inject your interferon to prevent skin irritation or a rash at the injection site. To manage gastrointestinal symptoms, eat small, frequent healthy meals or snacks rather than three large meals each day, and avoid spicy or fatty foods.



Many people find that side effects are worse when they first start taking a new drug but diminish over time. Talk to your health-care provider about other ways to manage treatment side effects, and inform him or her if your symptoms get worse.

11. HBV MANAGEMENT

Chronic hepatitis B can be a difficult disease to manage. However, there are steps you can take to cope with the disease and keep your liver as healthy as possible.

People infected with hepatitis B should receive regular medical attention. Biochemical liver tests should be monitored regularly (every 6-12 months). During treatment, levels of HBV DNA also should be measured to gauge how well therapy is working. People taking pegylated interferon should be monitored for side effects, including blood cell counts, thyroid function tests, and assessment of depression. People with inactive HBV should receive regular ALT tests. All HBV carriers should be screened for liver cancer. It is important to find a doctor who is knowledgeable about hepatitis B; hepatologists and gastroenterologists specialize in liver disease. If you are not comfortable with your doctor, ask family or friends to recommend someone they like.

12. HEALTHY LIVER TIPS

- Get regular health check-ups, including biochemical liver tests.
- Avoid or limit consumption of alcohol and recreational drugs.
- Take no more than the recommended doses of medications. Be careful when using multiple drugs, herbs, or drugs and herbs together.

- Eat a healthy, well-balanced diet.
- Get regular, moderate exercise.
- Inform all your health-care providers about all drugs, herbs, supplements, and alternative therapies you are using.
- Inform all your doctors and other health-care providers about your hepatitis B, especially if you need cancer treatment or steroid therapy.

Nutrition

Because the liver processes everything you eat and drink, a healthy, well-balanced diet is important. A healthy diet should follow the general guidelines based on ChooseMyPlate.gov – www.choosemyplate.gov. Such a diet is low in fat and sodium, high in complex carbohydrates, and has adequate protein. Today, dietary restrictions do not play a major role in managing chronic hepatitis B. Processed foods often contain chemical additives, so reduce consumption of canned, frozen, and other preserved foods. Eat organic fruits and vegetables to avoid pesticides and fertilizers. Read labels and become familiar with ingredients. Many doctors recommend that people with hepatitis avoid raw or undercooked shellfish such as oysters and clams, because they may contain infectious organisms or toxins. A well-balanced diet should contain all essential vitamins and minerals; avoid taking high-dose supplements – especially those containing vitamin A, vitamin D, or iron, which can be harmful to the liver. Consult a licensed dietitian for specific dietary recommendations and seek medical advice before undertaking any unconventional diet.

Alcohol, Drugs, and Toxins

People with hepatitis B – especially those with cirrhosis – should avoid consuming alcohol.

Certain drugs – prescription, over-the-counter, and recreational – and herbal remedies can be harmful to the liver (hepatotoxic), especially when taken in high doses or used in combination. People with HBV should inform their health-care providers about all drugs, herbal rem-

It is important to find a doctor who is knowledgeable about hepatitis B



edies, and supplements they are taking. Avoid or reduce consumption of recreational drugs. Do not exceed recommended drug doses.

Because the liver processes toxins, avoid exposure to toxic liquids and fumes such as solvents, paint thinners, pesticides, and aerosol sprays. If it is necessary to use such chemicals, work in a well-ventilated area, cover your skin, and wear gloves and a protective face mask.

General Wellness

Exercise – Regular aerobic exercise can improve overall fitness and may help reduce fatigue, stress, and depression. Most people with chronic hepatitis can safely engage in moderate exercise. Avoid exercise if you are feeling very ill. People with advanced cirrhosis should be cautious about lifting weights. All people with chronic hepatitis should consult their health-care provider before starting an exercise program.

Stress management – Controlling stress is a major factor in managing any chronic disease. Exercise, meditation, and time management can all help reduce stress. Try to maintain a realistic picture of your health and a positive attitude.

Managing fatigue – Fatigue and low energy levels are reported by some people with HBV. Learn your limits and try not to overextend yourself. Alternate strenuous activities with more restful ones. Take naps as needed and get enough sleep at night. Remember that your health is important; learn to say “no” to friends and family who have unrealistic expectations of your energy level.

Time management – Plan activities in advance and try to make realistic work and play schedules. Many people find it helpful to use a daily planner to help schedule and remember activities. Don’t forget to plan restful activities.

Meditation – Many people find meditation (a method of relaxation and clearing and focusing the mind) to be a useful tool in coping with chronic hepatitis. Meditation


can reduce stress and help you maintain a healthy outlook on life. There are many meditation traditions, some of which are simple and easy to learn.

Support Groups and Therapy

Many people with hepatitis B feel isolated and find it difficult to cope with the effects of living with a chronic illness. A support group can offer a safe space to discuss the emotional issues surrounding chronic hepatitis. In addition, information shared by peers can be helpful in making decisions, managing symptoms, and developing coping strategies. Some people also find therapy with a psychologist or social worker to be beneficial.

The Internet

The Internet contains a wealth of information – both good and bad. Always check the sources of the information you find online. Look for dates and references. Remember that not all the information you find on the Internet is correct. Talk to your doctor about any information you are concerned about. Common sense can go a long way! Visit the HCV Support Project web site at www.hbvadvocate.org and see the resource list for other recommended sites.



Most people with chronic hepatitis can safely engage in moderate exercise. Avoid exercise if you are feeling very ill.



13. CONCLUSION

While hepatitis B can have serious consequences, most people infected with HBV do not develop chronic disease and lead normal lives. Inactive carriers show evidence of HBV infection but do not have symptoms of liver dysfunction. A minority of people chronically infected with HBV develop progressive disease that can lead to serious liver damage, including cirrhosis, liver cancer, and liver failure. New drugs for HBV are being developed. Universal HBV vaccination presents the best hope for controlling the spread of hepatitis B. Beyond medical treatment, if you have HBV there are several steps you can take – including getting regular medical care, avoiding alcohol and drugs, eating a healthy diet, engaging in moderate exercise, managing stress and fatigue, and joining a support group – to keep your liver as healthy as possible and to improve your overall quality of life.

14. HEPATITIS RESOURCES

Web sites

CDC Hepatitis Branch home page: www.cdc.gov/

Hepatitis B Foundation: 215-489-4900; www.hepb.org

American Liver Foundation Hepatitis and Liver Disease Hotline: 800-223-0179; www.liverfoundation.org

American Association for the Study of Liver Disease – Guidelines for the Treatment of Chronic HBV in Adults and Children: <http://onlinelibrary.wiley.com/doi/10.1002/hep.28156/full>

Be Sure to Check Out These Other HBV Publications on our website:

Easy B Facts.

These fact sheets are one-page fact sheets that are simple and easy to understand on practically every topic related to hepatitis B. <http://hcvadvocate.org/hbv/easy-b-fact-sheets/>

HBV Facts.

These fact sheets are multi-page fact sheets on various issues that provide more in-depth information about Hepatitis B. <http://hcvadvocate.org/hbv/hbv-fact-series/>

This Guide was originally written by Liz Highleyman. It has been reviewed and updated by Alan Franciscus.