



HCSP FACT SHEET

• HCV AND WOMEN •

Pregnancy, Childbirth, and Breastfeeding

Written by: Lucinda K. Porter, RN

Foreword

Hepatitis C viral infection (HCV) is a medical problem that can create an emotional issue. It becomes even more emotional when we think that babies are at risk. The Centers for Disease Control and Prevention (CDC) estimate that the overall risk of mothers infecting their infants is about 4 to 7%. This rate increases when women also have HIV. The rate may be higher for those women who have a higher viral load (a higher amount of virus in their blood).

Pregnancy should not occur while a patient is being treated for HCV. Ribavirin, one of the drugs used for HCV treatment, has been linked to birth defects and miscarriages. There are strong warnings against this that apply both to female patients and female partners of male patients. Patients are told to use two reliable forms of birth control during HCV treatment and for 6 months after treatment has stopped.

There are no recommendations against pregnancy for women with HCV who are not on HCV treatment and whose partners are not on treatment. If an HCV positive woman does become pregnant, she should inform her medical team of her HCV status. There are no recommendations against amniocentesis. There are no recommendations regarding vaginal versus Cesarean section delivery. There may be a higher mother-to-infant transmission risk for those with both HCV and HIV. Some doctors believe that a Cesarean may decrease this risk, but there are no specific recommendations for this.

After the birth, the mother should tell her infant's pediatrician. The (CDC) recommends testing for children born to HCV+ mothers. Some babies may be born with the HCV antibody but they may actually lose this by the time they are 18 months old. It is advised to wait for 18 months after the birth before doing this test.

Pregnancy does not change HCV progression except if cirrhosis is present. Lab tests measuring liver function (liver enzymes) may fluctuate during pregnancy

HCSP FACT SHEET

*A publication of the
Hepatitis C Support Project*

**EXECUTIVE DIRECTOR,
EDITOR-IN-CHIEF,
HCSP PUBLICATIONS**
Alan Franciscus

DESIGN
*Leslie Hoex,
Blue Kangaroo Design*

PRODUCTION
C.D. Mazoff, PhD

CONTACT INFORMATION

Hepatitis C Support Project
PO Box 15144
Sacramento, CA 95813
alanfranciscus@hcvadvocate.org

The information in this fact sheet is designed to help you understand and manage HCV and is not intended as medical advice. All persons with HCV should consult a medical practitioner for diagnosis and treatment of HCV.

This information is provided by the Hepatitis C Support Project a nonprofit organization for HCV education, support and advocacy

Reprint permission is granted and encouraged with credit to the Hepatitis C Support Project.

Pregnancy, Childbirth, and Breastfeeding

and post-partum. The amount of virus in the blood (viral load) may also fluctuate during pregnancy. It is recommended that women needing a viral load test should wait until after the postpartum period.

Women should not breastfeed during HCV treatment. Otherwise, there are no recommendations against breastfeeding for HCV+ women. During this time, practice good nipple care. If your nipples are cracked or bleeding, stop breastfeeding until you are healed.

One question that HCV+ women ask is should they become pregnant first or have HCV treatment first. There is no clear answer to this. For those considering treatment, the overall success rate is around 80%. Women have a slightly higher successful outcome than

men. Those who are younger also have an advantage. Treatment lengths vary, but if your treatment duration is 48 weeks, that means you need to prevent pregnancy for at least 48 weeks plus another 6 months. Some people don't want to wait that long.

Women who start their family first might want to postpone treatment until their children are older. Current HCV treatment has side effects, such as fatigue and depression. It can be hard to take care of young children if you aren't in good physical and emotional condition. No matter what age your child or children are, get support while you are undergoing treatment. New HCV medications that are in the pipeline are expected to have fewer side effects and shorter treatment lengths.

Related publications:

- **HCV and Women: Women and HCV: Overview**
www.hcvadvocate.org/hepatitis/factsheets_pdf/Women_and_HCV.pdf
- **HCV and Women: Being a Positive Mother**
www.hcvadvocate.org/hepatitis/factsheets_pdf/Wm_Mother.pdf
- **Women and Hepatitis C: An HCSP Guide**
www.hcvadvocate.org/hepatitis/factsheets_pdf/Women_Guide.pdf

For more information

- | | |
|---|--|
| <ul style="list-style-type: none">• Centers for Disease Control and Prevention
www.cdc.gov• HealthFinder
www.healthfinder.gov | <ul style="list-style-type: none">• Help4Hep
(877) HELP4HEP (877) 435-7443
www.help4hep.org• Hepatitis C Support Project
www.hcvadvocate.org |
|---|--|

Visit our websites to learn more about viral hepatitis:

**www.hcvadvocate.org • www.hbvadvocate.org
www.hepatitistattoos.org**